

BAZHENOVA, K.M.; DEMIN, V.N.; STANKEVICH, A.A.

Second Leningrad Municipal Oncological Conference. Vop.onk. 5
(MIRA 12:12)

(ONCOLOGY--CONGRESSES)

STANKEVICH, A.A. (Loningrad, Lisiy Nos, Losinaya ul., d.18)

New-model cicrular colpostat for radium therapy. Vop.onk. 5 no.9:374-378 '59. (MIRA 12:12)


l. Iz kafedry onkologii (zav. - prof. A.I. Rakov) Gosudarstvennogo instituta dlya usovershenstvovaniya vrachey im. S.M. Kirova i radiye-voy laboratorii (zav. - doktor med.nauk N.D. Perumova) Instituta onko-logii AMN SSSR (dir. - deystvitel nyy chlen AMN SSSR prof. A.I. Serebrov).

(RADIUM ther.)


(CERVIX UTERINE neopl.)

```
STANKEVICH, A.A., kand.med.nauk (Leningrad, Lisiy Nos, Losinaya, d.18)
       Rectal complications in the treatment of cancer of the cervix
       with radiant energy. Vest.rent. i rad. 34 no.3:41-46 My-Je
       159.
                                                          (MIRA 12:10)
       1. Iz kafedry onkologii (zav. - prof.A.I.Rakov) Instituta dlya
       usovershenstvovaniya vrachey (dir. - prof.N.I.Blinov) i Instituta
       onkologii (dir. - chlen-korrespondent AMN SSSR prof.A.I.Serebrov)
       Akademii meditsinskikh nauk SSSR.
                  (CERVIX HEOPLASMS, ther.
                        radium & x-ray, rectal compl. (Rus))
                  (RADIUM, ther. use
                        cancer of cervix, rectal compl. (Rus))
                  (RADIOTHERAPY, in various dis.
                        same)
                   (RECTUM, dis.
                        caused by racium & x-ray ther. of cancer of
                        cervix (Rus))
```

Significance of secondary radiation from filters of Co⁶⁰ preparations during intracavitary use. Med.rad. 5 no.7:32-36 *60. (MIRA 13:12) (COBALT-ISOTOPES) (RADIATION-PHYSIOLOGICAL MFFEOT)



Protective safe for storing radioactive substances. Vop. onk. 6 no.4:79-81 Ap '60. (MIRA 14:3)



	STANKEV	ICH, A.	A.					17	
		Proteconk.	tive table 7 no.3:125-	for work -127 '61. (RADIA	with radi	oactive pr ECTION)	reparation	(MIRA 148	5)
								•	
÷									
1,2									
								€	

STANKEVICH, A.A. (Leningrad, Lisiy Nos. Losinaya ul., 18)

New construction of a protective safe for keeping radioactive substances. Vop. onk. 7 no. 4:117-120 161. (MIRA 14:4)

1. Iz radiyevoy laboratorii (zav. - doktor med. nauk N.D. Perumova) Instituta onkologii AMN SSSR (dir. - deystvitel'nyy chlen AMN SSSR prof. A.I. Serebrov). (RADIOACTIVITY—SAFETY MEASURES)

STANKEVICH, A. A.

Calculation of gamma ray doses from volumes of varying geometric form uniformly filled with the COOO radioisotope. Vop. onk. 8 no.5:80-87 62.

1. Iz radiyevoy laboratorii (zav. - d-r med. nauk N. D. Perumova) Instituta onkologii AMN SSSR (dir. - deystv. chl. AMN SSSR, prof. A. I. Serebrov)

> (RADIATION-DOSAGE) (COBALT-ISOTOPES)

STANKEVICH, A.A. (Leningrad, prospekt Engel'sa, 28, kv.111,

Determination of doses in intracavital treatment of cancer of the uterus taking into consideration the position of radio-active preparations in the polyis. Vop. onk. 8 nc.9175-79 162.

(MIRA 17:6)
1. Iz radiyevoy laboratorii (zav. doktor med. rauk N.D. Perumova)
Instituta onkologii &MN SOSR (dir. deystvitelinyy chlen AMN
SSSR, prof. A.T. Serebrov).

STANKEVICH, A.A.

Evaluation of the photographic method of dosimetry with small dimension cameras in radiotherapy of cancer of the cervix uteri. Vop. onl. 9 no.1:106-111 163. (MIRA 16:5)

1. Iz radiyevoy laboratorii (zav. doktor med.nauk N.D.Perumova)
Instituta onkologii AMN SSSR (direktor - deystvitel'nyy chlen
AMN SSSR Prof. A.I.Serebrov).

(UTERUS—CANCER) (PHOTOGRAPHY, MEDICAL)

(RADIATION—DOSAGE)

STANKEVION. A.A. (Leningrad, Belth, prospekt Engel'sa, 28. kv. 111)

Accelerated photographic method in establishing isodose curves in interstitial radioisotope therapy. Vop. onk. 9 no.10:103-108 163. (MIRA 17:12)

l. Fa radiyevoy laboratorii (zav. - doktor med. nauk N.D.Perumova) Instituta onkologii AMN SSSR (direktor - deystvitel'nyy chlen AMN SSSR prof. A.L.Serebrov).

STANKEVICH, A.A.

Calculation of doses in intracavitary transment of cancer of the uterus with radioactive preparations using a rectangular coordinate system. Vop. onk. 11 no.1:80-108 '65. (MIRA 18:6)

1. Iz radiyevoy laboratorii (zav. - doktor med.nauk N.D.Perumova)
Instituta onkologii AMN SSSR (dir. - deystvitel'nyy chlen AMN
SSSR prof. A.I.Serabrov).

CIA-RDP86-00513R001652830002-4 "APPROVED FOR RELEASE: 08/25/2000

STANKSVICH, A.A.

Economic efficiency of planued technological processes for core. making. Avt. prom. 31 no.5137-39 My 165.

(MIRA 18:5)

1. Minskiy filial Nauchno-issledovatel skogo instituta tekhnologii aviomobilinoy promyshlennosti.

STANKEVICH, A.A.

Calculation of doses in a plane not coincident with the plane of isodoses in intracavital treatment of cancer of the cervix uteri. (MIRA 18:6) Vop. onk. 11 no.3:106-113

1. Iz radiyevoy laboratorii (zav. - doktor med. nauk N.D. Perumova) Instituta onkologii AMN SSSR (dir. - deystvitel'nyy chlen AMN SSSR prof. A.I. Serebrov), Moskva.

20151

9.4300 (and 1147, 1155, 1158)

S/181/61/003/002/049/050 B102/B201

AUT HORS:

Smolenskiy, G. A., Chang Tsung, and Stankevich, A. K

TITLE:

Effect of electron diffusion upon the radio-frequency dispersion of the magnetic permeability of garnet-type

ferrites

PERIODICAL:

Fizika tverdogo tela, v. 3, no. 2, 1961, 663-667

TEXT: In weak electric and magnetic fields; certain ferrites display relaxation processes which are correlated with electron diffusion. The mechanism of these relaxation processes has never been fully clarified so far. In this connection, a study was made of the complex magnetic permeability and the complex dielectric constant, as well as of the dielectric and semiconductor properties (the latter were studied by Ya. M. Ksendzov and V. A. Stogova). Concerning the study of the dispersion of the magnetic permeability a report has already been given at the 3rd All-Union Conference concerned with physics, the physicochemical properties of ferrites, and the physical bases of their application (June, 1959. Minsk). The polycrystalline specimens were prepared by the usual ceramic technique, using analytically pure Card 1/5

20151

S/181/61/003/002/049/050 B102/B201

Effect of electron ...

Card 2/5


iron oxide and yttrium oxide consisting of Y203 for 99.9%. Initial and final sintering temperatures amounted to 1150 and 1450°C. The latter temperature was reduced by adding 0.5-1% copper oxide to the yttrium ferrite. Aside from the polycrystalline specimens also single crystals were prepared (by Titova) as well as the following solid solutions. Y3^{Fe}4.75^{Al}0.25^O12, Y3^{Fe}4^{AlO}12, and Y3^{Fe}4.8^{Cr}0.2^O12 Measurements included the frequency dependence of μ^+ and μ^- in weak fields $^{\hat{}}(H\sim 1\ millioersted)$ in the frequency range of 10 kc/sec-25 Mc/sec. The frequency dependence of μ^{\prime} and μ " of single crystals at room temperature is illustrated in Fig. 1. A study of the low-frequency maximum of μ^{μ} at different temperatures showed that it was to be identified as dispersion with relaxation mechanism. This maximum shifts toward a higher frequency with a rise of temperature. The mean activation energy was found to be U = 0.375 ev. Approximate calculations indicated that domain boundaries were displaced in the frequency and temperature ranges considered. The magnetic spectra of garnet-type ferrites remind one of the so-called double-dispersion spectra - shf dispersion arises beside r-f dispersion. The yttrium ferrites investigated were synthesized at high temperatures (1400-1560°C), and contained about 0.2% Fe2+ (of the total iron amount). Resistivity at room temperature ranged between

20151

S/181/61/003/002/049/050 B102/B201

Effect of electron ...

10⁶ and 10⁷ ohm.cm. When the specimens were heated in oxygen current, the concentration of Fe2+ ions was reduced, and resistivity increased. Fig. 2 shows the frequency dependence of μ ' and μ " at room temperature and H 2 1 moe of polycrystalline specimens prior to (curves 1 and 1') and after (2, 2') heating in oxygen current (15 hr at 1000°C). 1-2% of CuO was added to some of the specimens (curves 3 and 31), their resistivity ranged between 1010 and 1011 ohm cm at room temperature; similar results were obtained on specimens with 1-2% Mn₂O₃ addition (4, 4'). For a comparison, Fig. 2 shows, moreover, the frequency dependence of μ of single crystals (curve 5). single crystals had a resistivity of 1012 ohm.cm. A study of the three abovementioned solid solutions showed that μ ' is reduced with increasing Al³ concentration, and that the maximum of μ " is shifted toward higher frequencies. The introduction of Cr^{3+} increases μ^{1} . The magnetic and electric spectra (i.e., $\mu'(f)$ and $\mathcal{E}'(f)$) of the ferrites investigated have a similar course. In all cases where there arises electron diffusion, μ' and ϵ' attain high values at small frequencies. A final clarification of the effect of electron diffusion upon the dispersion of magnetic permeability requires further studies. V. A. Ioffe, A. G. Gurevich, and I. Ye. Gubler are men-Card 3/5



GRASHCHENKOV, N.I., professor; KASSIL', G.N. (Moskva): (Po materialam S.P. Vinitskovskoy, G.S. Vorsa, S.M. Grach, N.G. Grachenoy, M.B. Dunayevskoy F.A. Rosinoy, V.V. Stankevich. A.L. Sheakhmana, A.A. Shmidt)

Data on nasal reflex therapy in medical practice. Klin. med. 33 no. 9:12-17 S 155. (MIRA 9:2)

1. Iz terapavticheskogo, nervnogo i fizioterapevticheskogo otdeleniy Moskovskoy ordena Lenina bol'nitsy imeni S.P. Botkina i nauchnoissledovatel'skoy gruppy pri otdelenii biologicheskikh nauk Akademii nauk SSSR. 2. Deystvitel'nyy chlen AMN SSSR (for Grashchenkov) (THERAPEUTICS,

mass reflex ionogalvanic ther. technic)
(RLECTROTHERAPY,
mass reflex ionogalvanic ther. technic)

SHCHEGOLEV, Lev Illarionovich; EL'MANOVICH, Lidiya Yakovlevna; STANKEVICH, Anna L'vovna; YERMOLAYEVA, I.A., red.; LEBEDEVA, Z.V., tekhn. red.

[Textbook of the English language as an aid for reading and translating medical literature] Uchebnoe posobie po angliiskomu translating medical literature] Uchebnoe posobie po angliiskomu iazyku dlia chteniia i perevoda meditsinskoi literatury. Izd.2., ispr. i dop. Leningrad, Medgiz, 1962. 382 p. (MIRA 15:7) (ENGLISH LANGUAGE—TECHNICAL ENGLISH) (MEDICINE—TERMINOLOGY)

STANKEVICH, A.M.; STANKEVICH, I.M., inzh.

Measures which made possible the improvement of the technical conditions of NB-406 engines. Elek. i tepl.tiaga 6 no.8:16-18 (MIRA 17:3) Ag 162.

1. Zamestitel' nachal'nika depo Kurgan Yuzhno-Ural'skoy dorogi (for A.M.Stankevich). 2. Apparatnyy tsekh depo Kurgan Yuzhno-Ural'skoy dorogi (for I.M.Stankevich).

PODBEL'SKIY, G.N., kand.tekhn.mauk; STANKEVICH, A.S., inzh.

Industrial-genetic classification of humic coals (for discussion).
Nauch. trudy KuzNIIUgleobog. no.1:90-108 '62. (MIRA 16:8)

(Kuznetsk Basin--Coal--Classification)

BORCDULIN, V.A., inch.; STANKLVICH, A.S., inch.; ARTAMONOV, V.V., inch.

Investigating the effect of the depth of preparation on the coking properties of petrographic ally heterogenous Kuznetsk Basin coal. Nauch. trudy KuzNITUglechog. no.2:198-207 '64. (MIRA 17:10)

STANKSVICH, A.S., Inches ACCAMEMEN, V.V., Inches LUKANIN, A.A., Inches KORSHUNOV, V.A., Inches


Pilot plant coking of prepared coal from seams of lower subscries of the Balakorka series in the Prokop'yevsk-Kiselevsk region. Nauch. trudy KuzNIIUgleobog. no.2:207-212 164. (MIRA 17:10)

STANKEVICH, A.S., inzh.; PODBELISKIY, G.N., kand.tekhn.nauk

Using the method of the coking laboratory of the Institute of
Mineral Fuels to study the coking capacity of Kuznetsk Basin

coals. Nauch. trudy KuzNIIUgleobog. no.1:108-117 '62. (MIRA 16:8)


(Kuznetsk Basin--Coal--Carbonization)



KOLKER, O.N.; STANKEVICH, A.V.

Electronic automatic multiple-point chart-recording instruments and assemblies. Mash. i neft. obor. no.2:44-48 163. (MIRA 17:8)

1. Lenteplopribor.



STATEMENICE, F.S.: "The blood supply of the tendons of the thigh muscles."

Irkutsk State Medical Inst. Irkutsk, 1954.
(Dissertation for the Degree of Candidate in Medical Sciences)


So: Knizhnaya Letopis', No. 18, 1956


STANKEVICH, B.Ye.; ISAYEVA, M.I.

Selection of sites for air intake for ventilation of buildings at petroleum refineries. Gig. 1 san. no.6:27-34 Je '54. (MIRA 7:6)

1. Is Ufinekogo neftyanogo nauchno-issledovatel'ekogo instituta. (VENTILATION,

*selection of sites for air intake in petroleum-refining plants)





SOV/81-59-16-58505

Translation from: Referativnyy zhurnal. Khimiya, 1959, Nr 16, p 410 (USSR)

AUTHORS: Isayeva, M.I., Kalnina, R.V., Stankevich, B.Ye., Eygenson, A.S.

TITLE: The Alkalinization of Gasoline Distillates by Trisodiumphosphate

PERIODIC AL: Tr. Bashkirsk. n.-i. in-t po pererabotke nefti, 1959, Nr 1, pp 100-109

The results of the work of a pilot installation at the Ufa Oil Refinery ABSTRACT: are presented (a diagram is given). The gasoline distillate of thermal cracking at $44 - 200^{\circ}$ C with a H₂S content in the amount of 0.017 - 0.026 weight % after alkalinization with trisodiumphosphate (I) stands a test

with a copper plate. The recommended concentration of an aqueous I solution is 5 - 5.5 weight %, the sulfur content 7.5 g/l. The regeneration of the solution is carried out by boiling for 1 hour under vacuum at 120 - 130 mm Hg. On introducing alkalinization by I in oil refineries the consumption of NaOH and the quantity of sulfurous-alkaline industrial

sewage will decrease sharply. The purification of gasoline by I should

be cheaper than the purification by NaOH.

S. Rozenoyer.

Card 1/1

ISAYEVA, M.I.; STANKEVICH, B.Yo.; TOROPTSEV, N.G.

Ways for reducing caustic soda consumption in alkalizing clear petroleum products. Trudy BashNII NP no.1:110-119 '59.

(MIRA 12:6)

(Petroleum products)
(Sodium hydroxide)


STANKEVICH, B.Ye.; MITKALEV, B.A.; ISAYEVA, M.I.

Aeration purification of sewage containing hydrogen sulfide at petroleum refineries. Trudy BashMII MP no.1:205-215 '59.

(MIRA 12:6)

(Sewage-Purification) (Hydrogen sulfide)

(Petroleum refineries-By-products)



SOKOLAV, F.A.; STANKEVICH, B.Ye.; TOROPTSEV, N.G.

Development of methods for the utilization of sulfur removed in the refining of clear petroleum products. Khim.sera-i azotorg.soed.sod. v neft.i nefteprod. 3:407-410 *60. (MIRA 14:6)

STANKEVICH, B.Ye.

Developing the conditions for decalting sour Arlan oil. Trudy Wash NIINP no.5:22-32 462.

ST LAEVICH, E.A.

hesults of organizing compound treatment for patients with poliomyelitis sequelae in Kiev Province. Ortop., travm. i protez. 26 (MIRA 18:7)

1. Iz Ukrainskogo instituta ortopedii i travmatologii (dir. - dotsent I.P.Alekseyenko). Adres avtora: Kiyev 54, ul. Vorov-skogo, dom 27, Institut ortopedii i travmatologii.

5 (3) AUTHORS: Vanag, G. Ya., Gren, E. Ya.,

SOV/153-2-2-13/31

Stankevich, E. I.

TITLE:

Polycyclic Heterocyclic Compounds (Mnogoyadernyye geterotsiklicheskiye soyedineniya). I. 4-Phenyldibenzoylene

Pyridine (I. 4-fenil-dibenzoilenpiridin)

PERIODICAL:

Izvestiya vysshikh uchebnykh zavedeniy. Khimiya i khimicheskaya tekhnologiya, 1959, Vol 2, Nr 2, pp 210-214 (USSR)

ABSTRACT:

Recently the authors proved (Ref 1) that 4-phenyl-2,3 (CO), 6,5 (CO)-dibenzoylene pyridine (VII) develops when benzalindandione-1,3 (II) is heated with ammonium acetate in glacial acetic acid. The mechanism of this reaction was

explained. It proved that under the conditions of this reaction benzylindandione (II) partially decomposes in its primary compounds: benzaldehyde and indandione-1,3 (I). The latter immediately is added to the active ethylene linkage of the benzalindandione and forms diindandionylphenyl methane (III) (Ref 2). The compounds of the latter type easily split off a

water molecule from their enol forms (IV) and result in the corresponding pyranes (V). In these however the oxygen bridge is replaced by nitrogen under the influence of ammonia. In this

Card 1/3

Polycyclic Heterocyclic Compounds. I. 4-Phenyldi- SOV/153-2-2-13/31 benzoylene Pyridine

process they changed into the corresponding dihydropyridines (VI) (Refs 3-8). It proved however that heminal diindandione compounds of the type (III) are immediately changed into the corresponding 1,4-(or 3,4 ?)-dihydropyridines (Ref 9) under the influence of ammonium acetate. The mechanism of that change is not quite clear yet. These dihydropyridines are transformed into pyridines if exposed to the air (or quicker, if H2O2)(see scheme). Since the method given above is the quickest way for producing aryldibenzoylene pyridines, the problem should be dealt with in detail, in order to explain its scope of application. As expected, o- and p-nitrobenzal indandiones produced the corresponding nitrophenyl-dibenzoylene pyridines (VIII, IX) (Ref 2). In contrast to further statements given in reference 2, the authors succeeded in producing the corresponding dibenzoylene pyridines by means of heating the arylidene indandione which contained a nucleophilic substitute. The reaction however takes place much more slowly and the output is much lower. Anisal, salicylal, vanillal and veratral indandiones reacted positively

Card 2/3

Polycyclic Heterocyclic Compounds. I. 4-Phenyldibenzoylene Pyridine

SOV/153-2-2-13/31

in producing the substances X-XIII. All of the produced phenyldibenzoylene pyridines are yellow or orange substances with a very high (often over 300°) melting point, only the ortho-derivatives are crystalline. Their chemical activity is low. The rest of their properties is described. Since the acylates of the oxy compounds under discussion are yellow, and their alkaline salts are red or red violet, one has to draw the conclusion that during the salt production a tautomeric change of the oxy compounds takes place. Finally analogies of the recently published article, reference 10, are discussed. A simplification of the synthesis of the aryl-dibenzoylene pyridines can be attained, if the arylidene indandiones are not isolated. There are 13 references, 10 of which are Soviet.

ASSOCIATION:

Latviyskiy gosudarstvennyy universitet; Kafedra organicheskoy khimii (Latvian State University, Chair of Organic Chemistry)

SUBMITTED: Card 3/3

February 10, 1958

8/079/60/030/05/46/074 B005/B016

Vanag, G. Ya., Stankevich, E. I., Gren, E. Ya.

Polynuclear Heterocyclic Compounds, II. Structure and Color AUTHORS:

of Some Derivatives of 4-Phenyl-dibenzoylene Pyridine

PERIODICAL: Zhurnal obshchey khimii, 1960, Vol. 30, No. 5, pp. 1620-1627

TEXT: The authors of the present paper investigated the fine structure of 4-(p-dimethyl-amino-phenyl)-2,3(CO).6,5(CO)-dibenzoylene pyridine (II) and 4-(p-dimethyl-amino-m-nitro-phenyl)-2,3(CO).6,5(CO)-dibenzoyler. pyridine (III), as well as of arylidene indandiones which are the simplest representatives of this series. The absorption spectra of solutions of these compounds were taken in the ultraviolet and visible spectrum region and analyzed. Fig. 1 shows the ultraviolet absorption spectra of two phenyl-dibenzoylene pyridines in two different solvents (dioxane, C2H5OH + + C₂H₅ONa). Figs. 2 and 3 give the absorption spectra of solutions of

compound (II) in dioxane and in concentrated hydrochloric acid in the ultraviolet and visible spectrum region. For comparison, in each of these three

Card 1/2

DIANTTI

Edymonic Teneral delications, M. I., Creak S. T.,

Elymonical Chieffer S. T., Creak S. T.,

Frill This State of the property of the state of the sta

VANAG, G.Ya.; STANKEVICH, E.I.

APPROVED FOR RELEASE: 08/25/2000 CIA-RDP86-00513R001652830002-4" (dimedonyl) methanes with ammonium acetate. Zhur.ob.khim. 30 (MIRA 14:4)

1. Institut organicheskogo sinteza Akademii nauk Latviyskoy SSR. (Ammonium acetate) (Methane) (Acridinedione)

STANKEVICH, E.I .; VANAG, G.Ya. [Vanag*; G.], akademik

Asymmetric three-carbon condensations with 1,3-indandione. Dokl. AN SSSR 140 no.3:607-609 S '61. (MIRA 14:9)

1. Institut organicheskogo sinteza AN Latviyskoy SSR. 2. AN Latviyskoy SSR (for Vanag).

(Indandione) (Condensation products (Chemistry)

s/081/62/000/013/016/054 B158/B144

AUTHOR:

Stankevich, E.

TITLE:

Reaction of arylidenindandiones with imines of cyclic

β-diketones and ammonium acetate

FEFTODICAL: Referativnyy zhurnal. Khimiya, no. 13, 1962, 248, abstract

13Zn222 (Sb. "Tsiklich. β-diketony". Riga, AN LatvSSR,


1961, 269-274)

TEXT: Dimedone imine reacts with arylidenindandiones $Ar = C_6H_5$, -NO2C6H4-N(UH3)2C6H4, m-CH3O, p-C6H3OH] in CH3COOH; forming accordingly 2,3(CO)-benzoylene-7,7-dimethyl-4-aryl-5-keto-1,4,5,6,7,8-hexahydroquinolines, dark red compounds (a solution in alcoholic alkali is violetbiue) which are easily oxidized by HNO3 to the corresponding tetrahydroquinolines. An analogous reaction of benzal indandione with 5,5-dimethyl-3-butyl aminocyclohexene-2-one-1 leads to N-substituted benzoylene hexahydroquinoline. By heating arylidenirdandiones with ex-Card 1/2

Reaction of arylidenindandiones ...

\$/081/62/000/013/016/054 D:58/B144

cess CH3 COONH4 in glecial CH3 COOH, yellow or orange substances (Ia-m) are obtained. I is synthesized also by heating a mixture of indandione, aldenyde and CH3COONH, in glacial CH3COOH. The mechanism of the reaction is discussed as well as the formation of a dark colour on dissolving in alcoholic alkali. Data are given on uv-spectra for I.



In all cases $E = C_6H_4X$; (a) X = H, (b) $X = 2^{\dagger}-NO_2$, (c) $X = 3^{\dagger}-NO_2$,

(d) $\lambda = 4! = \text{NO}_2$, (e) $X = 4! - \text{N}(\text{CH}_3)_2$, (f) $X = 4! - \text{OCH}_3$, (g) $X = 3! - \text{OCH}_3$, $4! - \text{OH}_3$

(ii) X = 2! - OH, (i) $X = 3! - OCH_3$, $4! - OCH_3$, (k) X = 2! - OH, $3! - OCH_3$,

(1) $X = 5! - NO_2$, 4 $N(CH_3)_2$, (m) X = 3! - OH. [Abstracter's note: Complete translation. Card 2/2

ZHIZHEL', G.I., inzh.; STANKEVICH, E.M., inzh. Manufacture of pressureless socket pipes by centrifugation. Mekh. stroi. 19 no.4:14-16 Ap '62. (MIRA 15:9) (Pipe, Concrete)

VANAG, G.Ya.; STANKEVICH, E.Yu.; ROMADAN, Yu.P.

Improvement of the method for producing hexenal. Med.prom. 13 (MIRA 13:1) no.9:27-28 S 59.

1. Institut organicheskogo sinteza Akademii nauk Latviyskoy SSR. (HEXOBARBITAL)

Wontorer, L., insh.; STANKEVICH, F., inzh.

Using gas fuel in brick factories in the Ukraine. Stroi.mat. 4
no.10:24-26 0 '58.

(Ukraine --Gas as fuel) (Ukraine --Brickmaking)

BARANOV, L.A., inzh.; SKRYLEVA, G.I., inzh.; STANKEVICH, F.M.; VERTIKO., T.A.

Using alcohol-containing waste products from chemical industries as
a type of reagent in the flotation of coal slurry. Nauch.trudy KuzNITUglecbeg. no.2393-216 464.

DUEL', M.A., kand.tekhn.nauk; RABINOVICH, O.M., prof.; STANKEVICH, G.L., inzh.; FAYERSHTEYN, D.G., kand.tekhn.nauk

Testing the steam superheater of a high-pressure boiler fired with ash. Elek.sta. 29 no.8:22-25 Ag '58. (MIRA 11:11) (Superheaters-Testing)

STANKEVICH, G.L.

STANKEVICH, G.L.

"The Ukrainian S.S.R. and the Moldavian S.S.R.; economic map for secondary schools," Reviewed by G.L. Stankevich. Izv. Vses. geog. (MIRA 10:12) ob-va 89 no.6:561-563 K-D '57.

(Ukraine--Maps) (Moldavia--Maps) (Geography, Economic)

LIPOVETSKIY, S.Ye., inzh.; STANKEVICH, G.L., inzh.; FAYERSHTEYN, D.G., kand. tekhn. nauk

Utilizing the heat of the flue gases in burning natural gas under the steam boilers. Izv. vys. ucheb. zav.; energ. 2 no.7:69-73
J1 '59. (MIRA 13:1)

1. Khar'kovskiy politekhnicheskiy institut im. V.I. Lenina. (Boilers)

VYSOTSKAYA, A.I., inzh.; GORBATKO, P.A., inzh.; STANKEVICH, G.L., inzh.; FAYERSHTEYN, D.G., kand.tekhn.nauk

Complete analysis of blue gas in the combustion of natural gas under steam boilers. Izv.vys.ucheb.zav.; energ. 2 no.12: 85-89 D '59. (MIRA 13:5)


1. Khar'kovskiy politekhnicheskiy institut imeni V.I.Lenina Predstavlena kafedroy kotlostroyeniya. (Gas as fuel)

RABINOVICH, O.M., prof.; FAYERSHTEYE, D.G., kand.tekhn.nauk; STANKEVICH, G.L., inzh.; YEREMENKO, R.V. Testing a steam superheater of a boiler fired with natural gas. Elek.sta. 31 no.1:2-8 Ja '60. (MIRA 13:5) (Superheaters--Testing)

RABINOVICH, O.M., prof.; FAYERSHTEYN, D.G., kand.tekhn.nauk; STANKEVICH, G.L., inzh.

Experimental investigation of gas burners with peripheral gas feed. Elek.sta. 31 no.212-6 F '60. (MURA 13:5)

(Gas burners)



Gas burner with peripheral gas supply for large boiler units.

Elek. sta. 33 no.7:12-15 Jl '62.

(Boilers) (Gas burners)

KORBUG, Ye.V., inzh.; MERKHALEV, S.D., kand.tekhn.nauk; STANKEVICH, G.S., inzh.; Prinimal uchastiye PAVLOV, K.A.

Study of the discharge characteristics of soiled insulators. Elektrichestvo no.3:76-81 Mr 162. (MIRA 15:2)

1. Nauchno-issledovatel'skiy institut postoyannogo toka. (Electric insulators and insulation)

KORHUT, Ye.V.; MERKHALEV, S.D.; STANKEVICH, G.S.

Laboratory studies of the discharge characteristics of soiled insulators. Izv. NIIPT no.9:167-191 '62. (MIRA 15:12) (Electric lines—Overhead)

MERKHALEV, S.D., and tekhn nauk; STANKEVICH, G.S., inzh.

Duration of d.c. potential withstanding strength of suspension insulators during heavy rains. Elektrichestvo no.2:70-72 F '63. (MIRA 16:5)

Nauchno-issledovatel'skiy institut postoyannogo toka.
 (Electric power distribution) (Electric lines—Overhead)

STANKEVICH, I. A. Dr. Med. Sci.


Dissertation: "The Development of the Lower Sincipital Region in the Human." First Moscow Order of Lenin Medical Inst. 9 Jun 47.

SO: Vechernyaya Moskva, Jun, 1947 (Project #17836)

STANKEVICH, I.A.

Stankevich, I.A. "The development of the insular lobe of the human brain in the post-natal period", Trudy In-ta mozga (Gos. in-t mozga M-va zdravookhraneniya SSSR), Issue 6, 1948, p. 130-50, Tables XIX-XXII of an atlax (inserts).

SO: U-3042, 11 March 53, (Letopis 'nykh Statey, No. 9, 1949)



STANKEVICH, I.A., doktor med.nauk; NAUMOVA, T.S., kand.biol.nauk


Some recent data on the brain. Vest.AMN SSSR 11 no.5:24-34


(BRAIN


156.

morphol.structure & physical. funct., interrelation in health & dis., review.

(MIRA 12:10)







"Features of the structure of the human cerebrum and the temporal lobe of man and monkeys" by S.K.Blinkov. Reviewed by Ln.Stankewich, A.A.Khachaturian. Zbur.nevr. i psikh. 57 no.6:788-790 '57.

(BRAIN) (BLINKOV, S.M.)


(BLINKOV, S.M.)

KUZIMINA, A.V.; STANKEVICH, I.A.

"Trudy" of the Avicenna Medical Institute in Stalinbad, Papers of the Department of Normal Anatomy, vol. 14, no.1, 1955, vol.25, no.2, 1957. Reviewed by A.V. Kuz'mina, I.A. Stankevich. Arkh.anat. gist, i embr. 36 no.2:25-88 F 59. (MIRA 12:2)

1. Adres avtorov: Moskva, B-120, per. Obukha, d. 5, Institut mosga AMN SSSR.

(ANATOMY--PERICDICALS)




POPOVA, E.N., kand.biologicheskikh nauk; PREOBRAZHENSKAYA, N.S., doktor med.nauk; (STANKEVICH, I.A., doktor med.nauk

Results of a conference on the "Structure and function of the human analyzer in ontopeny." Vest. AMN SSSR 15 no.6:85-90 '60. (MIRA 14:4)

(BRAIN-LOCALIZATION OF FUNCTIONS)

APPROVED FOR RELEASE: 08/25/2000 CIA-RDP86-00513R001652830002-4"



SARKISOV, S.A., red.; KUKUYEV, L.A., red.; POLYAKOV, G.I., red.;

PREOBRAZHENSKAYA, N.S., red.; STANKEVICH, I.A., red.;

TROFIMOV, L.G., red.; ARKHANGEL'SKIY, Yu.V., red.; LYUDKOVSKAYA,

N.I., tekhn. red.

[Structure and function of the analysors of man in antogenesis] Struktura i funktsiia analizatorov cheloveka v ontogenezo; trudy. Pod obshchei red. S.A.Sarkisova. Moskva, Medgiz, 1961. (MIRA 15:12)

1. Rasshirennaya nauchnaya konferentsiya instituta mozga, 1959.
2. Deystvitel'nyy chlen Akademii meditsinskikh nauk SSSR (for Sarkisov). 3. Institut mozga Akademii medtisinskikh nauk SSSR, Moskva (for Polyakov, Kukuyev).

(SENSE-ORGANS) (ONTOGENY)

"Neuropathology and psychiatry" (Collected scientific works of neuropathologists and psychiatrists of the Latvian S. S. R.).

Reviewed by I.A. Stankevich and A.A. Khachaturian. Zhur. nevr. (All. (PSYCHIATRY))

(NERVOUS SYSTEM—DISEASES)

NAUMOVA, T.S.; STANKEVICH, I.A. (Moskva)


Results of the conference on the problem "Structure and function of the nervous system." Zhur. nevr. i psikh. 61 no.11:1737-1740 (NIRA 15'2)

'61. (NERVOUS SYSTEM)

STANKEVICH, I.A.

Comparative characteristics of the development of the cerebrum in man and monkeys. Zhur. nevr. i psikh. 61 no.12:1772-1780 (MIRA 15:7)

1. Institut mozga AMN SSSR, Moskva.
(BRAIN)



STANKEVICH, I.A.


CONTRACTOR SERVICE SER

"Problems in clinical neurology and psychiatry. Collected works of neuropathologists, neurosurgeons and psychiatrists in Estonia.

Vol. 1. Tallinn, 1961." Reviewed by I.A.Stankevich. Zhur.nerv.i psikh. 62 no.6:950-951 '62. (MIRA 15:11)

(ESTONIA -PSYCHIATRY)

(ESTONIA — PSYCHIATRY (ESTONIA — NEUROLOGY)




PREOBRAZHENSKAYA, N.S.; STANKEVICH, I.A.

Review of M.B. TSuker's back "Fundamentals of padiatric reuropathology." Zhur. news. i. psikh. 63 no.6:942-944 (63. (MIRA 17:6))

STANKEVICH, I. A.

"Sravnitel'naya kharakteristika onto i filogeneza bolyshogo mozga cheloveka i nizshey obez'yany."

reports submitted for 7th Intl Cong, Anthropological & Ethnological Sciences, Moscow, ,-10 Aug 64.



STANKEVICH, I.I.; NIKOLAYEV, A.F., prof., doktor tekhn. nauk, red.

[Graphic statics; a manual] Grafostatika; uchebnoe posobie.

[Moskva, Mosk. stankoinstrumental'nyy in-t, 1963. 38 p. (MIRA 17:7)

SHABEL NIKOV, G.P.; LISOVSKIY, G.D.; STANKEVICH, I.M.; RUDENKO, A.M.; LEDYAYKIN, S.D.; ZEMLYANOV, V.P.

Testing a system of sublevel caving with breaking and drawing of the ore in inclined layers. Gor. zhur. no.6:23-24

[MIRA 15:11]

1. Vsesoyuznyy nauchno-issledovatel'skiy institut tavetnykh metallov, Ust'-Kamenogorsk (for Shabel'nikov, Lisovskiy, Stankevich). 2. Salairskiy rudnik (for Rudenko, Ledyaykin, Zemlyanov).

(Salair region—Mining engineering)

APPROVED FOR RELEASE: 08/25/2000 CIA-RDP86-00513R001652830002-4"

STANKEVICH, A.M.: STANKEVICH, I.M., inzh.

Measures which made possible the improvement of the technical conditions of NB-406 engines. Elek. i tepl. tiaga 6 no.8:16-18

Ag '62. (MIRA 17:3)

1. Zamestitel' nachal'nika depo Kurgan Yuzhno-Ural'skoy dorogi (for A.M.Stankevich). 2. Apparatnyy tsekh depo Kurgan Yuzhno-Ural'skoy dorogi (for I.M.Stankevich).

CIA-RDP86-00513R001652830002-4 "APPROVED FOR RELEASE: 08/25/2000

SHKABARNYA, B.M., inzh.; SOLOV'YEV, G.A., inzh.; STANKEVICH, I.M., inzh.; LISOVSKIY, G.D., inzh.

Using reduced diameter boreholes. Gor. zhur. no.8:74 (MIRA 17:10) Ag 164.

1. Salairskiy rudnik (for Shkabarnya, Solov'yev).

2. Vsesoyuznyy nauchno-issledovatel skiy institut tsvetnoy metallurgii (for Stankevich, Lisovskiy).

STANKEVICH,

AUTHORS:

Bochvar, D. A., Stankevich, I. V., Chistjakov, A. L. 62-11-27/29

TITLE:

On the Relationship Between the Electron-Gas Method and the Molecular Orbit wethod (K scotnosheniyu mezhdu metodom elektron-

nogo gaza i metodom molekulyarnykh orbit)

PERIODICAL:

Izvestiya AN SSSR, Otdel.Khim.Nauk, 1957, Nr 11, pp. 1414-1414

ABSTRACT:

This is a letter to the editor. It is shown that instead of the usually applied formula:

 $+ \frac{2m}{2} E \Psi (x) = 0$ (1)

a much more common equation

 $+ Ak\Psi(x) = 0$

can be applied. (2)

That is to say, with the same boundary conditions, where A is a parameter, which is at our disposal. By this equation an oscillation system can easily combined, where a certain point $x(C_1)$ is opposed to the i.atom C. If the distance between the adjacent Catoms is equal, the p.coefficient of the j.linear combination of the molecular orbits methods becomes equal to the value of the j.equation (2) in the point $x(C_p)$. If the distance is different,

Card 1/2

SOV/62-58-6-31/37 Bochvar, D. A., Stankevich, I. V., AUTHORS: Chistyakov, A. b.

Letter to the Editor (Pis'ma redaktoru) Calculation of the Conjunction Energy in the \overline{S} -Triphenyl-Cyclopropenyl Cation TITLE:

(Raschet energii sopryazheniya dlya S-trifeniltsiklopropenil-

kationa)

Izvestiya Akademii nauk SSSR, Otdeleniye khimicheskikh nauk, PERIODICAL:

1958, Nr 6, pp. 793-793 (USSR)

In connection with the statement made concerning the synthesis of the S-triphenyl-cyclopropenyl cation (Ref 1) the salculation ABSTRACT:

of this compound was carried out by the LKAO MO-method in $\pi\text{-electron}$ approximation. The authors proceeded from the

following assumptions: 1) the o-skeleton is flat and shows the symmetry group C 3v)

2) all bond lengths are equal,

3) all Coulomb integrals are equal among themselves (equal to

4) all resonance integrals are equal (equal to β), 5) AO is passed over by overlapping integrals. Calculation Card 1/3

Letter to the Editor. Calculation of the SOV/62-58-6-31/37 Conjunction Energy in the \overline{S} -Triphenyl-Cyclopropenyl

Cation

showed that a closed electron shell (in the sense of Khykkel) exists. 20π -electrons of the system take up 10 molecular orbitals corresponding to their energy (in ascending order): $\alpha + 2.61 \beta$, $\alpha + 2.06 \beta$ (twofold degenerated level), $\alpha + 1.79 \beta$, $\alpha + 1.15 \beta$ (threefold degenerated level) and $\alpha + 0.76 \beta$. For the

 α + β (threefold degenerated level) and α + 0,76 β . For the compound discussed the conjunction energy (compared with the system of isolated binary bonds) is 9,16, β and exceeds the sum of the conjunction energies in phenyl rings and in the cyclopropenyl cation by 1,16 β . There is 1 reference.

្រ**ំ** ដោះប៉ែ

ASSOCIATION: Institut elementoorganicheskikh soyedineniy Akademii nauk SSSR

(Institute of Elemental-organisCompounds AS USSR)

SUBMITTED:

February 26, 1958

Card 2/3

Letter to the Editor . Calculation of the Conjunction Energy in the \bar{S} -Triphenyl-Cyclopropehyl Cation

SOV/62-58-6-31/37

1. Cyclic compounds--Properties 2. Cyclopropenyl ions--Energy 3. Mathematics

4. Perturbation theory

Card 3/3

5(4) AUTHORS: Bochvar, D. A., Gambaryan, N. P., Stankevich, I. V., Chistyakov, A. L. sov/76-32-12-22/32

TITLE:

A Qualitative Evaluation of the Stability of Heterocyclic Systems by Hueckel's Method of Approximation (O kachestvennoy otsenke ustoychivosti geterotsiklicheskikh sistem v ramkakh priblizheniya Gyukkelya)

PERIODICAL:

Zhurnal fizicheskoy khimii, 1958, Vol 32, Nr 12, pp 2797 - 2802 (USSR)

ABSTRACT:

E. Hueckel (Ref 1) used the words "closed electron shell" to explain the relative stability of cyclic ions. With molecules forming regular polygons of CH-groups, the first, not degenerate level is followed by several doubly degenerate levels. If ate level is followed by several doubly degenerate levels. If these levels are gradually filled in with π -electrons, closed electron shells are formed for systems with 2, 6, 10, 14 π -electrons in accordance with Pauli's principle. When a π -electrons in accordance with Pauli's principle. When a substitution takes place, the energy change may be considered a substitution takes place, the energy change may be considered as being a disturbance which does not exert any influence on the closed shell. A study is made of the general stability of

Card 1/2

CIA-RDP86-00513R001652830002-4 "APPROVED FOR RELEASE: 08/25/2000

24(5)

sov/56-36-2-48/63

AUTHORS:

Bochvar, D. A., Gambaryan, N. P., Stankevich, I. V.,

Chistyakov, A. L.

TITLE:

On Some Properties of Symmetry of the Eigenfunctions of the Equation of Schrödinger (O nekotorykh svoystvakh simmetrii

sobstvennykh funktsiy uravneniya Shredingera)

PERIODICAL:

Zhurnal eksperimental noy i teoreticheskoy fiziki, 1959,

Vol 36, Nr 2, pp 626-627 (USSR)

ABSTRACT:

The present paper deals with 2 facts hitherto (according to the authors' opinion) not discussed in literature. 1) The symmetry groups of the eigenfunctions of the Schrödinger (Shredinger) equation are subgroups of the symmetry group $G_{\widetilde{H}}$ of the corre-

statement 1) is sponding Hamiltonian H. 2) The contrary of not true, i.e. there are no subgroups of the group $G_{\mbox{\scriptsize H}}$ which are

not symmetry groups of the eigenfunctions of a given

Schrödinger equation. The proofs of the correctness of these 2 assertions are discussed step by step. The groups of the

solutions of a Schrödinger equation with a total system of eigenfunctions consist of all the possible co nels of the symmetry

Card i/2

SOV/56-36-2-48/63

On Some Properties of Symmetry of the Eigenfunctions of the Equation of Schrödinger

group of the Hamiltonian. There are 3 references, 1. of which is

Soviet.

ASSOCIATION: Institut elementoorganicheskikh soyedineniy Akademii nauk SSSR

(Institute of Element-Organic Compounds of the Academy of

Sciences, USSR)

SUBMITTED: October 25, 1958

Card 2/2

BOCHVAR, D.A.; STANKEVICH, I.V.; CHISTYAKOV, A.L.

THE PROPERTY OF THE PROPERTY O

Conjugation energies of the phenylcyclopropenyl and diphenylcyclopropenyl cations. Zhur. fiz. khim. 34 no. 11:2543-2545 N '60. (MIRA 14:1)

1. Akademiya nauk SSSR, Institut elementoorganicheskikh soyedineniy. (Cyclopropene) (Chemical bonds)

86827

s/020/60/135/005/015/043 B019/B067

24.4500

Bochvar, D. A., Stankevich, I. V., and Chistyakov, A. L.

TITLE:

AUTHORS:

Entropy of Localization and Extension in a Quantum Mechani-

cal System

PERIODICAL:

Doklady Akademii nauk SSSR, 1960, Vol. 135, No. 5,

pp. 1095-1096

TEXT: In a previous paper (Ref. 1), the authors together with N. P. Gambaryan suggested the definition of delocalization of a particle in a steady state of a quantum mechanical system as entropy of localization which might be calculated by appropriate eigenfunctions of the system. If $\psi(x_1, y_1, z_1, \dots, x_n, y_n, z_n)$ is the steady state of a system consisting of n particles, the probability density for the position of

the i-th particle is $\Phi(\tau_i) = \int |\gamma|^2 d\tau_1 \cdots d\tau_{i-1} d\tau_{i+1} \cdots d\tau_n, \text{ and the entropy}$ of the localization $h_i = -\int_{R_3} \Phi(\tau_i) \log \Phi(\tau_i) d\tau_i$. Here, R with the

Card 1/3

X

86827

Entropy of Localization and Extension in a Quantum Mechanical System

s/020/60/135/005/015/045 во19/во67

respective index denotes the space $d\tau_i = dx_i dy_i dz_i$, over which integration is made. In the present paper, a system is studied consisting of m + k particles. m particles (e.g., positive nuclei) are fixed in this system, k denotes the number of similar particles (e.g., electrons). The problem arises as to what degree this definition is connected with the concept of extension. The authors attempted to introduce a theoretical characteristic of extension into the quantum mechanical system considered here. They regard a coincidence of this quantum mechanical concept and the concept of space in the ordinary sense as necessary. It may then easily be demonstrated that with homogeneous distribution (constant density) in a given finite range D of the space R with a volume V_D (in the ordinary sense) the local entropy h which is determined by $h = -\int_D \rho \log_D \rho \, d\tau$ is $\log_D V_D$, i.e., $V_D = b^h$. In the following, the authors define the h-extension of particles in the quantum mechanical system (with given state) by $V_H = e^h$ volume units. It is found that the h-extension is independent of

Card 2/3

86827

Entropy of Localization and Extension in a Quantum Mechanical System

S/020/60/135/005/015/043 B019/B067

the base of the logarithm which proves the correctness of the definition. Finally, some examples are briefly discussed in which N. P. Gambaryan and E. S. Bogatova calculated the particle entropy in a potential well. There is ! Soviet reference.

PRESENTED:

June 29, 1960, by I. V. Obreimov, Academician.

SUBMITTED:

June 23, 1960

Card 5/3